INDEX TO AUTHORS

ALONSO, R. See DANDLIKER, HALBERT, FLORIN, ALONSO, and SCHAPIRO, 1029

ARCHER, GORDON T., and BLACKWOOD, ANGELA. Formation of Charcot-Leyden crystals in human eosinophils and basophils and study of the composition of isolated crystals, 173

ARQUILLA, EDWARD R., and FINN, JACK. Genetic control of combining sites of insulin antibodies produced by guinea pigs, 771

ATKINS, ELISHA, and HEIJN, CORNELIS, JR. Studies on tuberculin fever. III. Mechanisms involved in the release of endogenous pyrogen *in vitro*, 207

BALK, SAMUEL D. See PROSE, BALK, LIEBHABER, and KRUGMAN, 1151

BARNETT, SUSAN BELL. See JANDL, FILES, BARNETT, and MACDONALD, 299

BASSETT, EMMETT W., TANENBAUM, STUART W., PRYZWANSKY, KATHERINE, BEISER, SAM M., and KABAT, ELVIN A. Studies on human antibodies. III. Amino acid composition of four antibodies from one individual, 251

BAUM, STEPHEN G. See ROWE, and BAUM, 955

—. See ROWE, BAUM, PUGH, and HOGGAN, 943

BEAN, MICHAEL A. See JANOFF, SCHAEFER, SCHERER, and BEAN, 841

BEISER, SAM M. See BASSETT, TANENBAUM, PRYZWANSKY, BEISER, and KABAT, 251

BENDITT, EARL P. See NORMANN and BENDITT, 693, 709

BENSON, BELINDA. See COHN and BENSON, 455

BLACKWOOD, ANGELA. See ARCHER and BLACKWOOD, 173

BLEIWEIS, ARNOLD S., and KRAUSE, RICHARD M. The cell walls of Group D streptococci. I. The immunochemistry of the type 1 carbohydrate, 237

BOREL, YVES, FAUCONNET, MARTHE, and MIESCHER, PETER A. Effect of 6-mercaptopurine (6-MP) on different classes of antibody, 263

CHASE, RANDOLPH M., JR., and RAPAPORT, FELIX T. The bacterial induction of homograft sensitivity. I. Effects of sensitization with Group A streptococci, 721

—. See RAPAPORT and CHASE, 733

CHIAPPINO, GEROLAMO. See PERNIS, CHIAPPINO, KELUS, and GELL, 853

COCHRANE, CHARLES G., UNANUE, EMIL R., and DIXON, FRANK J. A role of polymorphonuclear leucocytes and complement in nephrotoxic nephritis, 99

—. See KNIKER and COCHRANE, 83

—. See WARD, COCHRANE, and MÜLLER-EBERHARD, 327

COHN, ZANVIL A., and BENSON, BELINDA. The *in vitro* differentiation of mononuclear phagocytes. III. The reversibility of granule and hydrolytic enzyme formation and the turnover of granule constituents, 455

COSTELLO, RICHARD. See DUBOS, COSTELLO, and SCHAEDLER, 1003

COSTELLO, RICHARD. See DUBOS, SCHAEDLER, COSTELLO, and HOET, 67
—. See SCHAEDLER, DUBOS, and COSTELLO, 59, 77

DAHL, LEWIS K., HEINE, MARTHA, and TASSINARI, LORRAINE. effects of chronic excess salt ingestion. Further demonstration that genetic factors influence the development of hypertension: evidence from experimental hypertension due to cortisone and to adrenal regeneration, 533
DANDLIKER, W. B., HALBERT, S. P., FLORIN, M. C., ALONSO, R., and SCHAPIRO, H. C. Study of penicillin antibodies by fluorescence polarization and immunodiffusion, 1029
DAVID, JOHN R. Suppression of delayed hypersensitivity *in vitro* by inhibition of protein synthesis, 1125
— and PATERSON, PHILIP Y. *In vitro* demonstration of cellular sensitivity in allergic encephalomyelitis, 1161
DIXON, FRANK J. See COCHRANE, UNANUE, and DIXON, 99
—. See UNANUE, LEE, DIXON, and FELDMAN, 565
DUBOS, RENÉ, COSTELLO, RICHARD, and SCHAEDLER, RUSSELL W. The influence of endotoxin administration on the nutritional requirements of mice, 1003
— SCHAEDLER, RUSSELL W., COSTELLO, RICHARD, and HOET, PHILIPPE. Indigenous, normal, and autochthonous flora of the gastrointestinal tract, 67
—. See MUSHIN and DUBOS, 745
—. See SCHAEDLER, DUBOS, and COSTELLO, 59, 77
DU BUY, HERMAN G., and JOHNSON, MARTIN L. Some properties of the lactic dehydrogenase agent of mice, 587
DUTTON, RICHARD W. Further studies of the stimulation of DNA synthesis in cultures of spleen cell suspensions by homologous cells in inbred strains of mice and rats, 759

EDELMAN, G. M. See MARCHALONIS and EDELMAN, 601
EVANS, R., and SALAMAN, M. H. Studies on the mechanism of action of Riley virus. III. Replication of Riley's plasma enzyme-elevating virus *in vitro*, 993

FAHEY, JOHN L., and SELL, STEWART. The immunoglobulins of mice. V. The metabolic (catabolic) properties of five immunoglobulin classes, 41
—. See TERRY, FAHEY, and STEINBERG, 1087
FAUCONNET, MARTHE. See BOREL, FAUCONNET, and MIESCHER, 263
FELDMAN, JOSEPH D. See UNANUE, LEE, DIXON, and FELDMAN, 565
FILES, NANCYE M. See JANDL, FILES, BARNETT, and MACDONALD, 299
FINN, JACK. See ARQUILLA and FINN, 771
FLORIN, M. C. See DANDLIKER, HALBERT, FLORIN, ALONSO, and SCHAPIRO, 1029
FRANGIONE, BLAS, and FRANKLIN, EDWARD C. Structural studies of human immunoglobulins. Differences in the Fd fragments of the heavy chains of G myeloma proteins, 1
FRANKLIN, EDWARD C. See FRANGIONE and FRANKLIN, 1
FREEDMAN, SAMUEL O. See GOLD and FREEDMAN, 467
FREI, EMIL, 3RD. See OPPENHEIM, WHANG, and FREI, 651

GELL, PHILIP G. H., and SELL, STEWART. Studies on rabbit lymphocytes *in vitro*. II. Induction

of blast transformation with antisera to six IgG allotypes and summation with mixtures of antisera to different allotypes, 813
GELL, PHILIP G. H. See PERNIS, CHIAPPINO, KELUS, and GELL,— 853
—. See SELL and GELL, 423, 923
—. See SELL, ROWE, and GELL, 823
GILES, JOAN P. See LIEBHABER, KRUGMAN, MCGREGOR, and GILES, 1135
GOEBEL, WALTHER F., and STAUB, ANNE MARIE. Colicine K. VI. The immune response of horses to a colicinogenic strain of *Escherichia coli*, 891
GOLD, PHIL, and FREEDMAN, SAMUEL O. Specific carcinoembryonic antigens of the human digestive system, 467
GOOD, ROBERT A. See HILGARD, MARTINEZ, and GOOD, 1017
GOWANS, J. L. See STROBER and GOWANS, 347
GREY, HOWARD M., and MANNIK, MART. Specificity of recombination of H and L chains from human γG-myeloma proteins, 619
GUSTAFSSON, BENGT E. See HAMMARSTRÖM, LAGERCRANTZ, PERLMANN, and GUSTAFSSON, 1075

HALBERT, S. P. See DANDLIKER, HALBERT, FLORIN, ALONSO, and SCHAPIRO, 1029
HAMMARSTRÖM, STEN, LAGERCRANTZ, RUTGER, PERLMANN, PETER, and GUSTAFSSON, BENGT E. Immunological studies in ulcerative colitis. II. Colon antigen and human blood group A- and H-like antigens in germfree rats, 1075
HARMON, CHARLES K. See ROTTA, PRENDERGAST, KARAKAWA, HARMON, and KRAUSE, 877
HEIJN, CORNELIS, JR. See ATKINS and HEIJN, 207
HEINE, MARTHA. See DAHL, HEINE, and TASSINARI, 533
HEROD, LLEWELLYN. See LANMAN and HEROD, 579
HEYSSEL, ROBERT M. See KOENIG, HEYSSEL, MELLY, and ROGERS, 117
HIJMANS, W. See VAN FURTH, SCHUIT, and HIJMANS, 1173
HILGARD, HENRY R., MARTINEZ, CARLOS, and GOOD, ROBERT A. Production of runt disease in tolerant mice by the injection of syngeneic lymphoid cells, 1017
HOET, PHILIPPE. See DUBOS, SCHAEDLER, COSTELLO, and HOET, 67
HOGGAN, M. DAVID. See ROWE, BAUM, PUGH, and HOGGAN, 943

ISAKOVIĆ, KATARINA, SMITH, STANLEY B., and WAKSMAN, BYRON H. Role of the thymus in tolerance. I. Tolerance to bovine γ-globulin in thymectomized, irradiated rats grafted with thymus from tolerant donors, 1103

JACKSON, ANNE L. See LANDY, SANDERSON, and JACKSON, 483
JANDL, JAMES H., FILES, NANCYE M., BARNETT, SUSAN BELL, and MACDONALD, RICHARD A. Proliferative response of the spleen and liver to hemolysis, 299
JANOFF, AARON, SCHAEFER, SONJA, SCHERER, JOAN, and BEAN, MICHAEL A. Mediators of inflammation in leucocyte lysosomes. II. Mechanism of action of lysosomal cationic protein upon vascular permeability in the rat, 841
JOHNSON, MARTIN L. See DU BUY and JOHNSON, 587
JONES, VALERIE E., and LESKOWITZ, SIDNEY. Immunochemical study of antigenic specificity in delayed hypersensitivity. IV. The production of unresponsiveness to delayed hypersensitivity with a single antigenic determinant, 505

JONSSON, NILS, and SJÖGREN, HANS O. Further studies on specific transplantation antigens in Rous sarcoma of mice, 403

KABAT, ELVIN A. See BASSETT, TANENBAUM, PRYZWANSKY, BEISER, and KABAT, 251

KARAKAWA, WALTER W., OSTERLAND, C. KIRK, and KRAUSE, RICHARD M. Detection of streptococcal group-specific antibodies in human sera, 195

—. See ROTTA, PRENDERGAST, KARAKAWA, HARMON, and KRAUSE, 877

KELUS, ANDREW S. See PERNIS, CHIAPPINO, KELUS, and GELL, 853

KENNY, KATHRYN. See ROBBINS, KENNY, and SUTER, 385

KNIKER, WILLIAM T., and COCHRANE, CHARLES G. Pathogenic factors in vascular lesions of experimental serum sickness, 83

KOENIG, M. GLENN, HEYSSEL, ROBERT M., MELLY, M. ANN, and ROGERS, DAVID E. The dynamics of reticuloendothelial blockade, 117

KRAUSE. See BLEIWEIS and KRAUSE, 237

—. See KARAKAWA, OSTERLAND, and KRAUSE, 195

—. See ROTTA, PRENDERGAST, KARAKAWA, HARMON, and KRAUSE, 877

KRUGMAN, SAUL. See LIEBHABER, KRUGMAN, MCGREGOR, and GILES, 1135

—. See PROSE, BALK, LIEBHABER, and KRUGMAN, 1151

KUNKEL, HENRY G. See MÅRTENSSON and KUNKEL, 799

LAGERCRANTZ, RUTGER. See HAMMARSTRÖM, LAGERCRANTZ, PERLMANN, and GUSTAFSSON, 1075

LANDY, MAURICE, SANDERSON, RONALD P., and JACKSON, ANNE L. Humoral and cellular aspects of the immune response to the somatic antigen of *Salmonella enteritidis*, 483

LANMAN, JONATHAN T., and HEROD, LLEWELLYN. Homograft immunity in pregnancy. The placental transfer of cytotoxic antibody in rabbits, 579

LEE, SUN. See UNANUE, LEE, DIXON, and FELDMAN, 565

LESKOWITZ, SIDNEY. See JONES and LESKOWITZ, 505

LIEBHABER, HARVEY, KRUGMAN, SAUL, MCGREGOR, DOROTHY, and GILES, JOAN P. Studies of a myxovirus recovered from patients with infectious hepatitis. I. Isolation and characterization, 1135

—. See PROSE, BALK, LIEBHABER, and KRUGMAN, 1151

MÅRTENSSON, LARS, and KUNKEL, HENRY G. Distribution among the γ-globulin molecules of different genetically determined antigenic specificities in the Gm system, 799

MACDONALD, RICHARD A. See JANDL, FILES, BARNETT, and MACDONALD, 299

MAHY, B. W. J., ROWSON, K. E. K., PARR, C. W., and SALAMAN, M. H. Studies on the mechanism of action of Riley virus. I. Action of substances affecting the reticuloendothelial system on plasma enzyme levels in mice, 967

—. See ROWSON, MAHY, and SALAMAN, 983

MALAWISTA, STEPHEN E. On the action of colchicine. The melanocyte model, 361

MANNIK, MART. See GREY and MANNIK, 619

MARCHALONIS, J., and EDELMAN, G. M. Phylogenetic origins of antibody structure. I. Multichain structure of immunoglobulins in

the smooth dogfish (*Mustelus canis*), 601
MARTINEZ, CARLOS. See HILGARD, MARTINEZ, and GOOD, 1017
MAURER, PAUL H. See PINCHUCK and MAURER, 665, 673
MCDEVITT, HUGH O., and SELA, MICHAEL. Genetic control of the antibody response. I. Demonstration of determinant-specific differences in response to synthetic polypeptide antigens in two strains of inbred mice, 517
MCGREGOR, DOROTHY. See LIEBHABER, KRUGMAN, MCGREGOR, and GILES, 1135
MELLORS, ROBERT C. Autoimmune disease in NZB/Bl mice. I. Pathology and pathogenesis of a model system of spontaneous glomerulonephritis, 25
MELLY, M. ANN. See KOENIG, HEYSSEL, MELLY, and ROGERS, 117
MIESCHER, PETER A. See BOREL, FAUCONNET, and MIESCHER, 263
MÖLLER, ERNA. Antagonistic effects of humoral isoantibodies on the *in vitro* cytotoxicity of immune lymphoid cells, 11
MÜLLER-EBERHARD, HANS J. See NILSSON and MÜLLER-EBERHARD, 277
—. See WARD, COCHRANE, and MÜLLER-EBERHARD, 327
MUSHIN, ROSE, and DUBOS, RENÉ. Colonization of the mouse intestine with *Escherichia coli*, 745

NILSSON, ULF R., and MÜLLER-EBERHARD, HANS J. Isolation of β_{1F}-globulin from human serum and its characterization as the fifth component of complement, 277
NOLAN, JAMES P., and O'CONNELL, CORNELIUS J. Vascular response in the isolated rat liver. I. Endotoxin, direct effects, 1063
NORMANN, SIGURD J., and BENDITT, EARL P. Function of the reticuloendothelial system. I. A study on the phenomenon of carbon clearance inhibition, 693
— and —. II. Participation of a serum factor in carbon clearance, 709

O'CONNELL, CORNELIUS J. See NOLAN and O'CONNELL, 1063
OPPENHEIM, JOOST J., WHANG, JACQUELINE, and FREI, EMIL, 3RD. The effect of skin homograft rejection on recipient and donor mixed leucocyte cultures, 651
OSOBA, DAVID. The effects of thymus and other lymphoid organs enclosed in Millipore diffusion chambers on neonatally thymectomized mice, 633
OSTERLAND, C. KIRK. See KARAKAWA, OSTERLAND, and KRAUSE, 195

PARR, C. W. See MAHY, ROWSON, PARR, and SALAMAN, 967
PATERSON, PHILIP Y. See DAVID and PATERSON, 1161
PERLMANN, PETER. See HAMMARSTRÖM, LAGERCRANTZ, PERLMANN, and GUSTAFSSON, 1075
PERNIS, BENVENUTO, CHIAPPINO, GEROLAMO, KELUS, ANDREW S., and GELL, PHILIP G. H. Cellular localization of immunoglobulins with different allotypic specificities in rabbit lymphoid tissues, 853
PINCHUCK, PAUL, and MAURER, PAUL H. Antigenicity of polypeptides (poly-α-amino acids). XV. Studies on the immunogenicity of synthetic polypeptides in mice, 665
— and —. XVI. Genetic control of immunogenicity of synthetic polypeptides in mice, 673

PRENDERGAST, THOMAS J. See ROTTA, PRENDERGAST, KARAKAWA, HARMON, and KRAUSE, 877
PRESSMAN, D. See ROHOLT, RADZIMSKI, and PRESSMAN, 785
PROSE, PHILIP H., BALK, SAMUEL D., LIEBHABER, HARVEY, and KRUGMAN, SAUL. Studies of a myxovirus recovered from patients with infectious hepatitis. II. Fine structure and electron microscopic demonstration of intracytoplasmic internal component and viral filament formation, 1151
PRYZWANSKY, KATHERINE. See BASSETT, TANENBAUM, PRYZWANSKY, BEISER, and KABAT, 251
PUGH, WENDELL E. See ROWE, BAUM, PUGH, and HOGGAN, 943

RADZIMSKI, G. See ROHOLT, RADZIMSKI, and PRESSMAN, 785
RAPAPORT, FELIX T., and CHASE, RANDOLPH M., JR. The bacterial induction of homograft sensitivity. II. Effects of sensitization with staphylococci and other microorganisms, 733
—. See CHASE and RAPAPORT, 721
RATNOFF, OSCAR D. Increased vascular permeability induced by human plasmin, 905
ROBBINS, JOHN B., KENNY, KATHRYN, and SUTER, EMANUEL. The isolation and biological activities of rabbit γM- and γG-anti-*Salmonella typhimurium* antibodies, 385
ROGERS, DAVID E. See KOENIG, HEYSSEL, MELLY, and ROGERS, 117
ROHOLT, O. A., RADZIMSKI, G., and PRESSMAN, D. Preferential recombination of antibody chains to form effective binding sites, 785
ROITT, I. M. See TORRIGIANI and ROITT, 181
ROTHBARD, SIDNEY, and WATSON, ROBERT F. Immunologic relations among various animal collagens, 441
ROTTA, JIRI, PRENDERGAST, THOMAS J., KARAKAWA, WALTER W., HARMON, CHARLES K., and KRAUSE, RICHARD M. Enhanced resistance to streptococcal infection induced in mice by cell wall mucopeptide, 877
ROWE, DAVID S. See SELL, ROWE, and GELL, 823
ROWE, WALLACE P., BAUM, STEPHEN G., PUGH, WENDELL E., and HOGGAN, M. DAVID. Studies of adenovirus SV40 hybrid viruses. I. Assay system and further evidence for hybridization, 943
— and —. II. Defectiveness of the hybrid particles, 955
ROWEN, ROBERT, and WIEST, MYRTLE A. A new acute phase protein elicited in mice by streptolysin O and other toxic or infectious agents, 547
ROWSON, K. E. K., MAHY, B. W. J., and SALAMAN, M. H. Studies on the mechanism of action of Riley virus. II. Action of substances affecting the reticuloendothelial system on the level of viremia, 983
—. See MAHY, ROWSON, PARR, and SALAMAN, 967

SALAMAN, M. H. See EVANS and SALAMAN, 993
—. See MAHY, ROWSON, PARR, and SALAMAN, 967
—. See ROWSON, MAHY, and SALAMAN, 983
SANDERSON, RONALD P. See LANDY, SANDERSON, and JACKSON, 483
SCHAEDLER, RUSSELL W., DUBOS, RENÉ, and COSTELLO, RICHARD. Association of germfree mice with bacteria isolated from normal mice, 77
—, —, and —. The development of

the bacterial flora in the gastrointestinal tract of mice, 59
SCHAEDLER, RUSSELL W. See DUBOS, COSTELLO, and SCHAEDLER, 1003
—. See DUBOS, SCHAEDLER, COSTELLO, and HOET, 67
SCHAEFER, SONJA. See JANOFF, SCHAEFER, SCHERER, and BEAN, 841
SCHAPIRO, H. C. See DANDLIKER, HALBERT, FLORIN, ALONSO, and SCHAPIRO, 1029
SCHERER, JOAN. See JANOFF, SCHAEFER, SCHERER, and BEAN, 841
SCHUIT, HENRICA R. E. See VAN FURTH, SCHUIT, and HIJMANS, 1173
SELA, MICHAEL. See MCDEVITT and SELA, 517
SELL, STEWART, and GELL, PHILIP G. H. Studies on rabbit lymphocytes *in vitro*. I. Stimulation of blast transformation with an antiallotype serum, 423
— and —. IV. Blast transformation of the lymphocytes from newborn rabbits induced by antiallotype serum to a paternal IgG allotype not present in the serum of the lymphocyte donors, 923
—, ROWE, DAVID S., and GELL, PHILIP G. H. Studies on rabbit lymphocytes *in vitro*. III. Protein, RNA, and DNA synthesis by lymphocyte cultures after stimulation with phytohemagglutinin, with staphylococcal filtrate, with antiallotype serum, and with heterologous antiserum to rabbit whole serum, 823
—. See FAHEY and SELL, 41
—. See GELL and SELL, 813
SIEGLER, RICHARD. Pathogenesis of thymic changes in NZB mice with hemolytic anemia, 929
SJÖGREN, HANS O. See JONSSON and SJÖGREN, 403
SMITH, STANLEY B. See ISAKOVIĆ, SMITH, and WAKSMAN, 1103
STASTNY, PETER, STEMBRIDGE, VERNIE A., VISCHER, THOMAS, and ZIFF, MORRIS. Homologous disease in the adult rat, a model for autoimmune disease. II. Findings in the joints, heart, and other tissues, 681
STAUB, ANNE MARIE. See GOEBEL and STAUB, 891
STEINBERG, ARTHUR G. See TERRY, FAHEY, and STEINBERG, 1087
STEMBRIDGE, VERNIE A. See STASTNY, STEMBRIDGE, VISCHER, and ZIFF, 681
STROBER, S., and GOWANS, J. L. The role of lymphocytes in the sensitization of rats to renal homografts, 347
SUTER, EMANUEL. See ROBBINS, KENNY, and SUTER, 385

TANENBAUM, STUART W. See BASSETT, TANENBAUM, PRYZWANSKY, BEISER, and KABAT, 251
TASSINARI, LORRAINE. See DAHL, HEINE, and TASSINARI, 533
TERRY, WILLIAM D., FAHEY, JOHN L., and STEINBERG, ARTHUR G. Gm and Inv factors in subclasses of human IgG, 1087
TORRIGIANI, G., and ROITT, I. M. The enhancement of 19S antibody production by particulate antigen, 181

UNANUE, EMIL R., LEE, SUN, DIXON, FRANK J., and FELDMAN, JOSEPH D. Experimental glomerulonephritis. VII. The absence of an autoimmune antikidney response in nephrotoxic serum nephritis, 565
—. See COCHRANE, UNANUE, and DIXON, 99

VAN FURTH, R., SCHUIT, HENRICA R. E., and HIJMANS, W. The immunological development of the human fetus, 1173

VISCHER, THOMAS. See STASTNY, STEMBRIDGE, VISCHER, and ZIFF, 681

WAKSMAN, BYRON H. See ISAKOVIĆ, SMITH, and WAKSMAN, 1103

WARD, PETER A., COCHRANE, CHARLES G., and MÜLLER-EBERHARD, HANS J. The role of serum complement in chemotaxis of leucocytes *in vitro*, 327

WATSON, ROBERT F. See ROTHBARD and WATSON, 441

WEIGLE, WILLIAM O. The production of thyroiditis and antibody following injection of unaltered thyroglobulin without adjuvant into rabbits previously stimulated with altered thyroglobulin, 1049

WHANG, JACQUELINE. See OPPENHEIM, WHANG, and FREI, 651

WIEST, MYRTLE A. See ROWEN and WIEST, 547

WILSON, DARCY B. Quantitative studies on the behavior of sensitized lymphocytes *in vitro*. I. Relationship of the degree of destruction of homologous target cells to the number of lymphocytes and to the time of contact in culture and consideration of the effects of isoimmune serum, 143

—. II. Inhibitory influence of the immune suppressor, imuran, on the destructive reaction of sensitized lymphoid cells against homologous target cells, 167

ZIFF, MORRIS. See STASTNY, STEMBRIDGE, VISCHER, and ZIFF, 681

INDEX TO SUBJECTS

ACID, amino, composition of four antibodies from one individual, 251
—, deoxyribonucleic, stimulation of synthesis in cultures of spleen cell suspensions by homologous cells in inbred strains, 759
—, ribonucleic and deoxyribonucleic, and protein synthesis by lymphocyte cultures after stimulation with phytohemagglutinin, staphylococcal filtrate, antiallotype serum, and heterologous antiserum to rabbit whole serum, 823
Acids, poly-α-amino, antigenicity of polypeptides, 665, 673
Adenovirus SV40 hybrid viruses, assay system and further evidence for hybridization, 943
— SV40 hybrid viruses, defectiveness of hybrid particles, 955
Adjuvant, unaltered thyroglobulin without, production of thyroiditis and antibody following injection into rabbits previously stimulated with altered thyroglobulin, 1049
Adrenal regeneration and cortisone, evidence from hypertension due to, genetic factors influencing development of hypertension, effects of chronic excess salt ingestion, 533
Allergic encephalomyelitis, *in vitro* demonstration of cellular sensitivity, 1161
Allotype, anti-, serum, phytohemagglutinin, staphylococcal filtrate, and heterologous antiserum to rabbit whole serum, after protein, RNA, and DNA synthesis by lymphocyte cultures, 823
Allotype, anti-, serum, stimulation of blast transformation with, lymphocytes *in vitro*, 423
—, anti-, serum to paternal IgG allotype not present in serum of lymphocyte donors, blast transformation of lymphocytes from newborn rabbits induced by, 923
—, IgG, paternal, not present in serum of lymphocyte donors, blast transformation of lymphocytes from newborn rabbits induced by anti-allotype serum, 923
— serum, anti-, phytohemagglutinin, staphylococcal filtrate, and heterologous antiserum to rabbit whole serum, after protein, RNA, and DNA synthesis by lymphocyte cultures, 823
Allotypes, six IgG, induction of blast transformation with antisera to, and summation with mixtures of antisera to different allotypes, lymphocytes *in vitro*, 813
Allotypic specificities in lymphoid tissues, different, cellular localization of immunoglobulins with, 853
Amino acid composition of four antibodies from one individual, 251
— acids, poly-α-, antigenicity of polypeptides, 665, 673
Anemia, hemolytic, pathogenesis of thymic changes in NZB mice with, 929
Antibodies, γM- and γG-anti-*Salmonella typhimurium*, isolation and biological activities, 385
—, human, amino acid composition of four antibodies from one individual, 251

Antibodies, insulin, produced by guinea pigs, genetic control of combining sites, 771
—, penicillin, study by fluorescence polarization and immunodiffusion, 1029
—, streptococcal group-specific, in human sera, detection, 195
Antibody chains, preferential recombination to form effective binding sites, 785
—, cytotoxic, placental transfer, homograft immunity in pregnancy, 579
—, effect of 6-mercaptopurine on different classes, 263
— production, 19S, enhancement by particulate antigen, 181
— response, genetic control of, demonstration of determinant-specific differences in response to synthetic polypeptide antigens in two strains of inbred mice, 517
— structure, phylogenetic origins, multichain structure of immunoglobulins in *Mustelus canis*, 601
— and thyroiditis, production following injection of unaltered thyroglobulin without adjuvant into rabbits previously stimulated with altered thyroglobulin, 1049
Antigen, colon, and human blood group A- and H-like antigens in germfree rats, immunological studies in ulcerative colitis, 1075
—, particulate, enhancement of 19S antibody production, 181
—, somatic, of *Salmonella enteritidis*, immune response to, humoral and cellular aspects, 483
Antigenic specificities in Gm system, distribution among γ-globulin molecules of different genetically determined, 799
—, specificity in delayed hypersensitivity, immunochemical study, production of unresponsiveness, to delayed hypersensitivity with single antigenic determinant, 505
Antigenicity of polypeptides, poly-α-amino acids, 665, 673
Antigens, carcinoembryonic, of human digestive system, 467
—, human blood group A- and H-like, and colon antigen in germfree rats, immunological studies in ulcerative colitis, 1075
—, specific transplantation, in Rous sarcoma, 403
—, synthetic polypeptide, in two strains of inbred mice, demonstration of determinant-specific differences in response to, genetic control of antibody response, 517
Antisera to six IgG allotypes, induction of blast transformation with, and summation with mixtures of antisera to different allotypes, lymphocytes *in vitro*, 813
Antiserum to rabbit whole serum, heterologous, phytohemagglutinin, staphylococcal filtrate, and anti-allotype serum, after protein, RNA, and DNA synthesis by lymphocyte cultures, 823
Autoimmune antikidney response in nephrotoxic serum nephritis, absence, glomerulonephritis, 565
— disease in NZB/Bl mice, pathology and pathogenesis of model system of spontaneous glomerulonephritis, 25
— disease, model for, homologous disease in adult rat, findings in joints, heart, and other tissues, 681

BACTERIA isolated from normal mice, association with germfree mice, 77
Bacterial flora, development in gastrointestinal tract, 59
— induction of homograft sensitivity, effects of sensitization with Group A streptococci, 721
— induction of homograft sensitivity, effects of sensitization with staphylococci and other microorganisms, 733

Basophils and eosinophils, human, formation of Charcot-Leyden crystals in, and study of composition of isolated crystals, 173
Biological activities and isolation of γM- and γG-anti-*Salmonella typhimurium* antibodies, 385
Blood, human, group A- and H-like antigens and colon antigen in germfree rats, immunological studies in ulcerative colitis, 1075
Bovine γ-globulin in thymectomized, irradiated rats grafted with thymus from tolerant donors, tolerance, 1103

CARBOHYDRATE, type 1, immunochemistry, cell walls of Group D streptococci, 237
Carbon clearance inhibition, phenomenon, function of reticuloendothelial system, 693
— clearance, participation of serum factor in, function of reticuloendothelial system, 709
Carcinoembryonic antigens of human digestive system, 467
Catabolic, metabolic, properties of five immunoglobulin classes, 41
Cationic protein, lysosomal, mechanism of action upon vascular permeability, mediators of inflammation in leucocyte lysosomes, 841
Cell suspensions, spleen, stimulation by homologous cells in inbred strains, DNA synthesis in cultures of, 759
— wall mucopeptide, enhanced resistance to streptococcal infection induced in mice by, 877
— walls of Group D streptococci, immunochemistry of type 1 carbohydrate, 237
Cells, homologous target, relationship of degree of destruction to number of lymphocytes and time of contact in culture and consideration of effects of isoimmune serum, 143
Cells, immune lymphoid, *in vitro* cytotoxicity of, antagonistic effects of humoral isoantibodies, 11
—, sensitized lymphoid, destructive reaction against homologous target cells, inhibitory influence of immune suppressor, imuran, on, 167
—, syngeneic lymphoid, production of runt disease in tolerant mice by injection, 1017
Cellular and humoral aspects of immune response to somatic antigen of *Salmonella enteritidis*, 483
— localization of immunoglobulins with different allotypic specificities in lymphoid tissues, 853
— sensitivity in allergic encephalomyelitis, *in vitro* demonstration, 1161
Chain, multichain, structure of immunoglobulins in *Mustelus canis*, phylogenetic origins of antibody structure, 601
Chains, antibody, preferential recombination to form effective binding sites, 785
—, H and L, from human γG-myeloma proteins, specificity of recombination, 619
—, heavy, of G myeloma proteins, differences in Fd fragments, structural studies of human immunoglobulins, 1
Charcot-Leyden crystals, formation in human eosinophils and basophils, and study of composition of isolated crystals, 173
Chemical, immuno-, study of antigenic specificity in delayed hypersensitivity, production of unresponsiveness to delayed hypersensitivity with single antigenic determinant, 505
Chemistry, immuno-, of type 1 carbohydrate, cell walls of Group D streptococci, 237
Chemotaxis of leucocytes *in vitro*, role of serum complement, 327
Colchicine, melanocyte model, 361

Colicine K, immune response of horses to colicinogenic strain of *Escherichia coli*, 891
Colicinogenic strain of *Escherichia coli*, immune response of horses to, 891
Colitis, ulcerative, immunological studies in, colon antigen and human blood group A- and H-like antigens in germfree rats, 1075
Collagens, immunologic relations among, 441
Colon antigen and human blood group A- and H-like antigens in germfree rats, immunological studies in ulcerative colitis, 1075
Complement, fifth component of, isolation from human serum, characterization of β_{1F}-globulin as, 277
— and polymorphonuclear leucocytes in nephrotoxic nephritis, 99
—, serum, in chemotaxis of leucocytes *in vitro*, 327
Cortisone and adrenal regeneration, evidence from hypertension due to, genetic factors influencing development of hypertension, effects of chronic excess salt ingestion, 533
Crystals, Charcot-Leyden, formation in human eosinophils and basophils and study of composition of isolated crystals, 173
Culture, time of contact in, and number of lymphocytes, relationship of degree of destruction of homologous target cells to, and consideration of effects of isoimmune serum, 143
Cultures, DNA synthesis in, spleen cell suspensions, stimulation by homologous cells in inbred strains, 759
—, lymphocyte, protein, RNA, and DNA synthesis by, after stimulation with phytohemagglutinin, staphylococcal filtrate, antiallotype serum, and heterologous antiserum to rabbit whole serum, 823
Cultures, recipient and donor mixed leucocyte, effect of skin homograft rejection on, 651
Cytotoxic antibody, placental transfer, homograft immunity in pregnancy, 579
Cytotoxicity, *in vitro*, of immune lymphoid cells, antagonistic effects of humoral isoantibodies, 11

DEHYDROGENASE agent, lactic, properties, 587
Deoxyribonucleic acid, protein, and RNA synthesis by lymphocyte cultures after stimulation with phytohemagglutinin, staphylococcal filtrate, antiallotype serum, and heterologous antiserum to rabbit whole serum, 823
— acid synthesis, stimulation in cultures of spleen cell suspensions by homologous cells in inbred strains, 759
Diffusion chambers, Millipore, thymus and lymphoid organs enclosed in, effects on neonatally thymectomized mice, 633
—, immuno-, and fluorescence polarization, study of penicillin antibodies by, 1029
Digestive system, human, carcinoembryonic antigens, 467
Disease, autoimmune, in NZB/Bl mice, pathology and pathogenesis of model system of spontaneous glomerulonephritis, 25
—, homologous, in adult rat, model for autoimmune disease, findings in joints, heart, and tissues, 681
—, runt, production in tolerant mice by injection of syngeneic lymphoid cells, 1017

ELECTRON microscopic and fine structure demonstration of intracytoplasmic internal component and viral filament formation, myxovirus recovered from patients with infectious hepatitis, 1151

Encephalomyelitis, allergic, *in vitro* demonstration of cellular sensitivity, 1161
Endogenous pyrogen *in vitro*, mechanisms involved in release, tuberculin fever, 207
Endotoxin administration, influence on nutritional requirements, 1003
—, direct effects, vascular response in isolated rat liver, 1063
Enzyme-elevating virus, Riley's plasma, replication of *in vitro*, mechanism of action of Riley virus, 993
—, hydrolytic, and granule formation, reversibility of, and turnover of granule constituents, differentiation of mononuclear phagocytes, 455
— levels, plasma, action of substances affecting reticuloendothelial system on, mechanism of action of Riley virus, 967
Eosinophils and basophils, human, formation of Charcot-Leyden crystals in, and study of composition of isolated crystals, 173
Escherichia coli, colicinogenic strain, immune response of horses to, 891
— *coli*, colonization of intestine, 745

FETUS, human, immunological development, 1173
Fever, tuberculin, mechanisms involved in release of endogenous pyrogen *in vitro*, 207
Filament formation, viral, and intracytoplasmic internal component, fine structure and electron microscopic demonstration of, myxovirus recovered from patients with infectious hepatitis, 1151
Filtrate, staphylococcal, phytohemagglutinin in, antiallotype serum, and heterologous antiserum to rabbit whole serum, after protein, RNA, and DNA synthesis by lymphocyte cultures, 823
Flora, bacterial, development in gastrointestinal tract, 59
—, indigenous, normal, and autochthonous, of gastrointestinal tract, 67
Fluorescence polarization and immunodiffusion, study of penicillin antibodies by, 1029

GASTROINTESTINAL tract, development of bacterial flora in, 59
— tract, indigenous, normal, and autochthonous flora, 67
Genetic control of antibody response, demonstration of determinant-specific differences in response to synthetic polypeptide antigens in two strains of inbred mice, 517
— control of combining sites of insulin antibodies produced by guinea pigs, 771
— control of immunogenicity of synthetic polypeptides, 673
— factors influencing development of hypertension, evidence from hypertension due to cortisone and adrenal regeneration, effects of chronic excess salt ingestion, 533
Genetically determined antigenic specificities in Gm system, distribution among γ-globulin molecules of, 799
Germfree mice, association with bacteria isolated from normal mice, 77
— rats, colon antigen and human blood group A- and H-like antigens in, immunological studies in ulcerative colitis, 1075
Globulin, bovine γ-, tolerance in thymectomized, irradiated rats grafted with thymus from tolerant donors, 1103
— G, immuno-, allotype, paternal, not present in serum of lymphocyte donors, blast transformation of lymphocytes from newborn rabbits induced by antiallotype serum to, 923

Globulin G, immuno-, allotypes, six, induction of blast transformation with antisera to, and summation with mixtures of antisera to different allotypes, lymphocytes *in vitro*, 813
— G, immuno-, subclasses of human, Gm and Inv factors, 1087
—, immuno-, five classes, metabolic, catabolic, properties, 41
—, β_{1F}-, isolated from human serum and characterization as fifth component of complement, 277
— molecules, γ-, of different genetically determined antigenic specificities in Gm system, distribution, 799
Globulins, immuno-, human, structural studies, differences in Fd fragments of heavy chains of G myeloma proteins, 1
—, immuno-, in *Mustelus canis*, multichain structure, phylogenetic origins of antibody structure, 601
—, immuno-, with different allotypic specificities in lymphoid tissues, cellular localization, 853
Glomerulonephritis, abscence of autoimmune anitkidney response in nephrotoxic serum nephritis, 565
—, spontaneous, pathology and pathogenesis of model system, autoimmune disease in NZB/Bl mice, 25
Grafted, thymectomized, irradiated rats, with thymus from tolerant donors, tolerance to bovine γ-globulin, 1103
Granule and hydrolytic enzyme formation and turnover of granule constituents, reversibility of, *in vitro* differentiation of mononuclear phagocytes, 455

HEART, joints, and tissues, findings in, homologous disease in adult rat, model for autoimmune disease, 681
Hemolysis, proliferative response of spleen and liver to, 299
Hemolytic anemia, pathogenesis of thymic changes in NZB mice with, 929
Hepatitis, infectious, myxovirus recovered from patients with, fine structure and electron microscopic demonstration of intracytoplasmic internal component and viral filament formation, 1151
—, infectious, myxovirus recovered from patients with, isolation and characterization, 1135
Heterologous antiserum to rabbit whole serum, phytohemagglutinin, staphylococcal filtrate, and antiallotype serum, after protein, RNA, and DNA synthesis by lymphocyte cultures, 823
Homograft immunity in pregnancy, placental transfer of cytotoxic antibody, 579
— rejection, skin, effect on recipient and donor mixed leucocyte cultures, 651
— sensitivity, bacterial induction of, effects of sensitization with Group A streptococci, 721
— sensitivity, bacterial induction of, effects of sensitization with staphylococci and other microorganisms, 733
Homografts, renal, lymphocytes in sensitization of rats to, 347
Homologous cells in inbred strains, stimulation of DNA synthesis in cultures of spleen cell suspensions, 759
— disease in adult rat, model for autoimmune disease, findings in joints, heart, and tissues, 681
— target cells, destructive reaction of sensitized lymphoid cells against, inhibitory influence of immune suppressor, imuran, 167
— target cells, relationship of degree of destruction to number of lymphocytes and time of contact in

culture, consideration of effects of isoimmune serum, 143
Humoral and cellular aspects of immune response to somatic antigen of *Salmonella enteritidis*, 483
— isoantibodies, antagonistic effects on *in vitro* cytotoxicity of immune lymphoid cells, 11
Hybrid viruses, adenovirus SV40, assay system and further evidence for hybridization, 943
— viruses, adenovirus SV40, defectiveness of hybrid particles, 955
Hybridization, adenovirus SV40 hybrid viruses, assay system and further evidence for, 943
Hydrolytic enzyme and granule formation and turnover of granule constituents, reversibility of, differentiation of mononuclear phagocytes, 455
Hypersensitivity, delayed, immunochemical study of antigenic specificity in, production of unresponsiveness to delayed hypersensitivity with single antigenic determinant, 505
— *in vitro*, delayed, suppression by inhibition of protein synthesis, 1125
Hypertension, genetic factors influencing development, evidence from hypertension due to cortisone and adrenal regeneration, effects of chronic excess salt ingestion, 533

IMMUNE lymphoid cells, *in vitro* cytotoxicity of, antagonistic effects of humoral isoantibodies, 11
— response of horses to colicinogenic strain of *Escherichia coli*, 891
— response to somatic antigen of *Salmonella enteritidis*, humoral and cellular aspects, 483
— suppressor, imuran, inhibitory influence on destructive reaction of sensitized lymphoid cells against homologous target cells, 167
Immunity, homograft, in pregnancy, placental transfer of cytotoxic antibody, 579
Immunochemical study of antigenic specificity in delayed hypersensitivity, production of unresponsiveness to delayed hypersensitivity with single antigenic determinant, 505
Immunochemistry of type 1 carbohydrate, cell walls of Group D streptococci, 237
Immunodiffusion and fluorescence polarization, study of penicillin antibodies by, 1029
Immunogenicity of synthetic polypeptides, 665
— of synthetic polypeptides, genetic control, 673
Immunoglobulin classes, five, metabolic, catabolic, properties, 41
— G allotype, paternal, not present in serum of lymphocyte donors, blast transformation of lymphocytes from newborn rabbits induced by antiallotype serum to, 923
— G allotypes, six, induction of blast transformation with antisera to, and summation with mixtures of antisera to different allotypes, lymphocytes *in vitro*, 813
— G, subclasses of human, Gm and Inv factors, 1087
Immunoglobulins, human, structural studies, differences in Fd fragments of heavy chains of G myeloma proteins, 1
— in *Mustelus canis*, multichain structure, phylogenetic origins of antibody structure, 601
— with different allotypic specificities in lymphoid tissues, cellular localization, 853
Immunologic relations among collagens, 441
Immunological development of human fetus, 1173
— studies in ulcerative colitis, colon antigen and human blood group

A- and H-like antigens in germfree rats, 1075
Imuran, immune suppressor, inhibitory influence on destructive reaction of sensitized lymphoid cells against homologous target cells, 167
Inbred mice, two strains of, demonstration of determinant-specific differences in response to synthetic polypeptide antigens in, genetic control of antibody response, 517
— strains, stimulation of DNA synthesis in cultures of spleen cell suspensions by homologous cells, 759
Infection, streptococcal, induced in mice by cell wall mucopeptide, enhanced resistance to, 877
Infectious hepatitis, myxovirus recovered from patients with, fine structure and electron microscopic demonstration of intracytoplasmic internal component and viral filament formation, 1151
— hepatitis, myxovirus recovered from patients with, isolation and characterization, 1135
— or other toxic agents and streptolysin O, new acute phase protein from elicitation by, 547
Inflammation in leucocyte lysosomes, mediators of, mechanism of action of lysosomal cationic protein upon vascular permeability, 841
Ingestion, salt, effects of chronic excess, further demonstration that genetic factors influence development of hypertension, evidence from experimental hypertension due to cortisone and adrenal regeneration, 533
Inhibition, carbon clearance, phenomenon, function of reticuloendothelial system, 693
— of protein synthesis, suppression of delayed hypersensitivity *in vitro*, 1125
Inhibitory influence of immune suppressor, imuran, on destructive reaction of sensitized lymphoid cells against homologous target cells, 167
Injection of syngeneic lymphoid cells, production of runt disease in tolerant mice, 1017
— of unaltered thyroglobulin without adjuvant into rabbits previously stimulated with altered thyroglobulin, production of thyroiditis and antibody following, 1049
Insulin antibodies produced by guinea pigs, genetic control of combining sites, 771
Intestine, colonization with *Escherichia coli*, 745
Intracytoplasmic internal component and viral filament formation, fine structure and electron microscopic demonstration, myxovirus recovered from patients with infectious hepatitis, 1151
Irradiated, thymectomized rats grafted with thymus from tolerant donors, tolerance to bovine γ-globulin in, 1103
Isoantibodies, humoral, antagonistic effects on *in vitro* cytotoxicity of immune lymphoid cells, 11
Isoimmune serum, consideration of effects, and relationship of degree of destruction of homologous target cells to number of lymphocytes and to time of contact in culture, 143
Isolated crystals, formation of Charcot-Leyden crystals in human eosinophils and basophils and study of composition, 173
—, rat liver, direct effects of vascular response with endotoxin, 1063
Isolation and characterization of myxovirus recovered from patients with infectious hepatitis, 1135

JOINTS, heart, and tissues, findings, homologous disease in adult rat, model for autoimmune disease, 681

KIDNEY, anti-, response in nephrotoxic serum nephritis, absence of autoimmune, glomerulonephritis, 565
—, homografts, lymphocytes in sensitization of rats to, 347

LACTIC dehydrogenase agent, properties, 587
Lesions, vascular, of serum sickness, pathogenic factors, 83
Leucocyte cultures, recipient and donor mixed, effect of skin homograft rejection on, 651
— lysosomes, mediators of inflammation, mechanism of action of lysosomal cationic protein upon vascular permeability, 841
Leucocytes *in vitro*, chemotaxis of, role of serum complement, 327
—, polymorphonuclear, and complement in nephrotoxic nephritis, 99
Liver, isolated rat, direct effects of vascular response with endotoxin, 1063
— and spleen, proliferative response to hemolysis, 299
Lymphocyte cultures, protein, RNA, and DNA synthesis by, after stimulation with phytohemagglutinin, staphylococcal filtrate, antiallotype serum, and heterologous antiserum to rabbit whole serum, 823
— donors, serum, blast transformation of lymphocytes from newborn rabbits induced by antiallotype serum to paternal IgG allotype not present in, 923
Lymphocytes from newborn rabbits, blast transformation, induced by antiallotype serum to paternal IgG allotype not present in serum of lymphocyte donors, 923
— in sensitization of rats to renal homografts, 347
—, in vitro, induction of blast transformation with antisera to six IgG allotypes and summation with mixtures of antisera to different allotypes, 813
Lymphocytes *in vitro*, protein, RNA, and DNA synthesis by lymphocyte cultures after stimulation with phytohemagglutinin, staphylococcal filtrate, antiallotype serum, and heterologous antiserum to rabbit whole serum, 823
— *in vitro*, stimulation of blast transformation with antiallotype serum, 423
—, number of, and time of contact in culture, relationship of degree of destruction of homologous target cells to, and consideration of effects of isoimmune serum, 143
—, sensitized, quantitative studies on behavior *in vitro*, 143, 167
Lymphoid cells, immune, *in vitro* cytotoxicity of, antagonistic effects of humoral isoantibodies, 11
— cells, sensitized, destructive reaction against homologous target cells, inhibitory influence of immune suppressor, imuran, 167
— cells, syngeneic, production of runt disease in tolerant mice by injection, 1017
— organs, thymus, enclosed in Millipore diffusion chambers, effects on neonatally thymectomized mice, 633
— tissues, cellular localization of immunoglobulins with different allotypic specificities, 853
Lysosomal cationic protein, mechanism of action upon vascular permeability, mediators of inflammation in leucocyte lysosomes, 841
Lysosomes, leucocyte, mediators of inflammation, mechanism of action of lysosomal cationic protein upon vascular permeability, 841

MAN, antibodies in, amino acid composition of four antibodies from one individual, 251
—, blood group A- and H-like antigens of, and colon antigen in germfree

rats, immunological studies in ulcerative colitis, 1075
Man, digestive system of, carcinoembryonic antigens in, 467
—, eosinophils and basophils in, formation of Charcot-Leyden crystals in, and study of composition of isolated crystals, 173
—, fetus of, immunological development, 1173
—, γG-myeloma proteins in, specificity of recombination of H and L chains, 619
—, IgG in, Gm and Inv factors in subclasses of, 1087
—, immunoglobulins in, structural studies, differences in Fd fragments of heavy chains of G myeloma proteins, 1
—, plasmin in, increased vascular permeability induced by, 905
—, sera in, streptococcal group-specific antibodies in, detection, 195
—, serum in, isolation of β_{1F}-globulin from, characterization as fifth component of complement, 277
Melanocyte model, colchicine, 361
6-Mercaptopurine, effect on different classes of antibody, 263
Metabolic, catabolic, properties of five immunoglobulin classes, 41
Microorganisms, staphylococci, effects of sensitization with, bacterial induction of homograft sensitivity, 733
Microsccopic, electron, and fine structure, demonstration of intracytoplasmic internal component and viral filament formation, myxovirus recovered from patients with infectious hepatitis, 1151
Millipore diffusion chambers, thymus and other lymphoid organs enclosed in, effects on neonatally thymectomized mice, 633
Molecules, γ-globulin, of different genetically determined antigenic specificities in Gm system, distribution, 799
Mononuclear phagocytes, *in vitro* differentiation, reversibility of granule and hydrolytic enzyme formation and turnover of granule constituents, 455
Mucopeptide, cell wall, enhanced resistance to streptococcal infection induced in mice by, 877
Mustelus canis, multichain structure of immunoglobulins in, phylogenetic origins of antibody structure, 601
Myeloma, human γG-, proteins, specificity of recombination of H and L chains, 619
— proteins, G, heavy chains of, differences in Fd fragments, structural studies of human immunoglobulins, 1
Myxovirus recovered from patients with infectious hepatitis, fine structure and electron microscopic demonstration of intracytoplasmic internal component and viral filament formation, 1151
— recovered from patients with infectious hepatitis, isolation and characterization, 1135

NEONATALLY thymectomized mice, effects of thymus and lymphoid organs enclosed in Millipore diffusion chambers on, 633
Nephritis, glomerulo-, spontaneous, pathology and pathogenesis of model system, autoimmune disease in NZB/Bl mice, 25
—, nephrotoxic, polymorphonuclear leucocytes and complement in, 99
—, nephrotoxic serum, absence of autoimmune antikidney response in, glomerulonephritis, 565
Nephrotoxic nephritis, polymorphonuclear leucocytes and complement in, 99
—, serum nephritis, absence of auto-

immune antikidney response in, glomerulonephritis, 565
Newborn rabbits, lymphocytes, induced by antiallotype serum to a paternal IgG allotype not present in serum of lymphocyte donors, blast transformation, 923
Nucleic acid, deoxyribo-, stimulation of synthesis in cultures of spleen cell suspensions by homologous cells in inbred strains, 759
— acid, ribo- and deoxyribo-, and protein synthesis by lymphocyte cultures after stimulation with phytohemagglutinin, staphylococcal filtrate, antiallotype serum, and heterologous antiserum to rabbit whole serum, 823
Nutritional requirements, influence of endotoxin administration, 1003

ORGANS, thymus and lymphoid enclosed in Millipore diffusion chambers, effects on neonatally thymectomized mice, 633

PARTICLES, hybrid, adenovirus SV40 hybrid viruses, defectiveness, 955
Pathogenesis of thymic changes in NZB mice with hemolytic anemia, 929
— and pathology of model system of spontaneous glomerulonephritis, autoimmune disease in NZB/Bl mice, 25
Pathogenic factors in vascular lesions of serum sickness, 83
Pathology and pathogenesis of model system of spontaneous glomerulonephritis, autoimmune disease in NZB/Bl mice, 25
Penicillin antibodies, study by fluorescence polarization and immunodiffusion, 1029
Permeability, increased vascular, induced by human plasmin, 905
—, vascular, mechanism of action of lysosomal cationic protein upon, mediators of inflammation in leucocyte lysosomes, 841
Phagocytes, mononuclear, *in vitro* differentiation, reversibility of granule and hydrolytic enzyme formation and turnover of granule constituents, 455
Phylogenetic origins of antibody structure, multichain structure of immunoglobulins in *Mustelus canis*, 601
Phytohemagglutinin, staphylococcal filtrate, antiallotype serum, heterologous antiserum to rabbit whole serum, after protein, RNA, and DNA synthesis by lymphocyte cultures, 823
Placental transfer of cytotoxic antibody, homograft immunity in pregnancy, 579
Plasma enzyme - elevating virus, Riley's, replication *in vitro*, mechanism of action of Riley virus, 993
-- enzyme, levels, action of substances affecting reticuloendothelial system on, mechanism of action of Riley virus, 967
Plasmin, human, increased vascular permeability induced by, 905
Polarization, fluorescence, and immunodiffusion, study of penicillin antibodies, 1029
Polymorphonuclear leucocytes and complement in nephrotoxic nephritis, 99
Polypeptide antigens, synthetic, in two strains of inbred mice, demonstration of determinant-specific differences in response, genetic control of antibody response, 517
Polypeptides, poly-α-amino acids, antigenicity, 665, 673
—, synthetic, genetic control of immunogenicity, 673
—, synthetic, immunogenicity, 665
Pregnancy, homograft immunity in, placental transfer of cytotoxic antibody, 579

Protein, lysosomal cationic, mechanism of action upon vascular permeability, mediators of inflammation in leucocyte lysosomes, 841
—, new acute phase, elicited by streptolysin O and other toxic or infectious agents, 547
—, RNA, and DNA synthesis by lymphocytes cultures after stimulation with phytohemagglutinin, staphylococcal filtrate, antiallotype serum, and heterologous antiserum to rabbit whole serum, 823
— synthesis, suppression of delayed hypersensitivity *in vitro* by inhibition, 1125
Proteins, G myeloma, heavy chains, differences in Fd fragments, structural studies of human immunoglobulins, 1
—, human γG-myeloma, specificity of recombination of H and L chains, 619
Purine, 6-mercapto-, effect on different classes of antibody, 263
Pyrogen, *in vitro*, endogenous, mechanisms involved in release, tuberculin fever, 207

REACTION, destructive, of sensitized lymphoid cells against homologous target cells, inhibitory influence of immune suppressor, imuran, on, 167
Recombination of H and L chains from human γG-myeloma proteins, specificity, 619
—, preferential, of antibody chains to form effective binding sites, 785
Regeneration, adrenal, and cortisone, evidence from hypertension due to, genetic factors influencing development of hypertension, effects of chronic excess salt ingestion, 533
Renal. *See* Kidney.
Resistance, enhanced, to streptococcal infection induced in mice by cell wall mucopeptide, 877
Reticuloendothelial blockade, dynamics, 117
— system, action of affecting substances on level of viremia, mechanism of action of Riley virus, 983
— system, action of affecting substances on plasma enzyme levels, mechanism of action of Riley virus, 967
— system, function, participation of serum factor in carbon clearance, 709
— system, function, phenomenon of carbon clearance inhibition, 693
Ribonucleic acid, deoxy-, protein, and RNA synthesis by lymphocyte cultures after stimulation with phytohemagglutinin, staphylococcal filtrate, antiallotype serum, and heterologous antiserum to rabbit whole serum, 823
— acid, deoxy-, stimulation of synthesis in cultures of spleen cell suspensions by homologous cells in inbred strains, 759
— acid, protein, and DNA synthesis by lymphocyte cultures after stimulation with phytohemagglutinin, staphylococcal filtrate, antiallotype serum, and heterologous antiserum to rabbit whole serum, 823
Riley virus, mechanism of action of substances affecting reticuloendothelial system on level of viremia, 983
— virus, mechanism of action of substances affecting reticuloendothelial system on plasma enzyme levels, 967
— virus, mechanism of action, replication of Riley's plasma enzyme-elevating virus *in vitro*, 993
Rous sarcoma, specific transplantation antigens, 403
Runt disease, production in tolerant mice by injection of syngeneic lymphoid cells, 1017

SALMONELLA enteritidis, immune response to somatic antigen of, humoral and cellular aspects, 483
— *typhimurium* antibodies, γM- and γG-anti–, isolation and biological activities, 385
Salt ingestion, effects of chronic excess, further demonstration that genetic factors influence development of hypertension, evidence from experimental hypertension due to cortisone and adrenal regeneration, 533
Sarcoma, Rous, specific transplantation antigens, 403
Sensitivity, cellular, in allergic encephalomyelitis, *in vitro* demonstration, 1161
—, homograft, bacterial induction of, effects of sensitization with Group A streptococci, 721
—, homograft, bacterial induction of, effects of sensitization with staphylococci and other microorganisms, 733
Sensitization of rats to renal homografts, role of lymphocytes in, 347
— with Group A streptococci, effects, bacterial induction of homograft sensitivity, 721
— with staphylococci and other microorganisms, effects, bacterial induction of homograft sensitivity, 733
Sensitized lymphocytes *in vitro*, quantitative studies on, 143, 167
— lymphoid cells against homologous target cells, inhibitory influence of immune suppressor, imuran, on destructive reaction of, 167
Sera, human, streptococcal group-specific antibodies in, detection, 195
Serum, antiallotype, phytohemagglutinin, staphylococcal filtrate, and heterologous antiserum to rabbit whole serum, after protein, RNA, and DNA synthesis by lymphocyte cultures, 823
Serum, antiallotype, stimulation of blast transformation with, lymphocytes *in vitro*, 423
—, antiallotype, to paternal IgG allotype not present in serum of lymphocyte donors, blast transformation of lymphocytes from newborn rabbits induced by, 923
— complement in chemotaxis of leucocytes *in vitro*, 327
— factor, participation in carbon clearance, function of reticuloendothelial system, 709
—, human, isolation of β_{1F}-globulin from, its characterization as fifth component of complement, 277
—, isoimmune, consideration of effects and relationship of degree of destruction of homologous target cells to number of lymphocytes and to time of contact in culture, 143
— nephritis, nephrotoxic, absence of autoimmune antikidney response in, glomerulonephritis, 365
—, rabbit whole, heterologous antiserum to phytohemagglutinin, staphylococcal filtrate, and antiallotype serum, after protein, RNA, and DNA synthesis by lymphocyte cultures, 823
— sickness, pathogenic factors in vascular lesions, 83
Skin homograft rejection, effect on recipient and donor mixed leucocyte cultures, 651
Somatic antigen of *Salmonella enteritidis*, immune response to, humoral and cellular aspects, 483
Specific, determinant-, differences in response to synthetic polypeptide antigens in two strains of inbred mice, demonstration, genetic control of antibody response, 517
—, group-, streptococcal antibodies in human sera, detection, 195

Specificities, antigenic, in Gm system, distribution among the γ-globulin molecules of different genetically determined, 799
—, different allotypic, in lymphoid tissues, cellular localization of immunoglobulins with, 853
Specificity, antigenic, in delayed hypersensitivity, immunochemical study, production of unresponsiveness to delayed hypersensitivity with single antigenic determinant, 505
— of recombination of H and L chains from human γG-myeloma proteins, 619
Spleen cell suspensions, DNA synthesis in cultures of, stimulation by homologous cells in inbred strains, 759
— and liver, proliferative response to hemolysis, 299
Staphylococcal filtrate, phytohemagglutinin, antiallotype serum, heterologous antiserum to rabbit whole serum, after protein, RNA, and DNA synthesis by lymphocyte cultures, 823
Staphylococci and other microorganisms, effects of sensitization with, bacterial induction of homograft sensitivity, 733
Stimulated, rabbits previously, with altered thyroglobulin, production of thyroiditis and antibody following injection of unaltered thyroglobulin without adjuvant, 1049
Stimulation of blast transformation with antiallotype serum, lymphocytes *in vitro*, 423
— of DNA synthesis in cultures of spleen cell suspensions by homologous cells in inbred strains of mice and rats, 759
— with phytohemagglutinin, staphylococcal filtrate, antiallotype serum, and heterologous antiserum to rabbit whole serum, after protein, RNA, and DNA synthesis by lymphocyte cultures, 823
Streptococcal group-specific antibodies in human sera, detection, 195
Streptococcal infection induced in mice by cell wall mucopeptide, enhanced resistance to, 877
Streptococci, Group A, effects of sensitization with, bacterial induction of homograft sensitivity, 721
—, Group D, cell walls of, immunochemistry of type 1 carbohydrate, 237
Streptolysin O and other toxic or infectious agents, new acute phase protein from elicitation by, 547
Syngeneic lymphoid cells, production of runt disease in tolerant mice by injection, 1017
Synthesis, DNA, stimulation in cultures of spleen cell suspensions by homologous cells in inbred strains, 759
—, protein, RNA, and DNA, by lymphocyte cultures after stimulation with phytohemagglutinin, staphylococcal filtrate, antiallotype serum, and heterologous antiserum to rabbit whole serum, 823
—, protein, suppression of delayed hypersensitivity *in vitro* by inhibition, 1125
Synthetic polypeptide antigens in two strains of inbred mice, demonstration of determinant-specific differences in response, genetic control of antibody response, 517
— polypeptides, genetic control of immunogenicity, 673
— polypeptides, immunogenicity, 665

THYMECTOMIZED, irradiated rats grafted with thymus from tolerant donors, tolerance to bovine γ-globulin in, 1103
— mice, neonatally, effects of thymus and other lymphoid organs enclosed in Millipore diffusion chambers on, 633
Thymic changes in NZB mice with

hemolytic anemia, pathogenesis, 929
Thymus in tolerance, role, 1103
— and other lymphoid organs enenclosed in Millipore diffusion chambers, effects on neonatally thymectomized mice, 633
Thyroglobulin, unaltered, injection without adjuvant into rabbits previously stimulated with altered thyroglobulin, production of thyroiditis and antibody following, 1049
Thyroiditis and antibody, production following injection of unaltered thyroglobulin without adjuvant into rabbits previously stimulated with altered thyroglobulin, 1049
Tissues, joints, heart, homologous disease in adult rat, model for autoimmune disease, 681
—, lymphoid, cellular localization of immunoglobulins with different allotypic specificities in, 853
Tolerance, role of thymus, 1103
— to bovine γ-globulin in thymectomized, irradiated rats grafted with thymus from tolerant donors, 1103
Tolerant mice, production of runt disease by the injection of syngeneic lymphoid cells, 1017
Toxic or infectious agents and streptolysin O, new acute phase protein from elicitation by, 547
Transformation, blast, induction with antisera to six IgG allotypes, and summation with mixtures of antisera to different allotypes, lymphocytes *in vitro*, 813
—, blast, of lymphocytes from newborn rabbits induced by antiallotype serum to paternal IgG allotype not present in serum of lymphocyte donors, 923
—, blast, stimulated by antiallotype serum, lymphocytes *in vitro*, 423
Transplantation, specific, antigens in Rous sarcoma, 403
Tuberculin fever, mechanisms involved in release of endogenous pyrogen *in vitro*, 207

ULCERATIVE colitis, immunological studies in, colon antigen and human blood group A- and H-like antigens in germfree rats, 1075

VASCULAR lesions of serum sickness, pathogenic factors, 83
— permeability induced by human plasmin, increased, 905
— permeability, mechanism of action of lysosomal cationic protein upon, mediators of inflammation in leucocyte lysosomes, 841
— response in isolated rat liver, endotoxin, direct effects, 1063
Viral filament formation and intracytoplasmic internal component, fine structure and electron microscopic demonstration, myxovirus recovered from patients with infectious hepatitis, 1151
Viremia, action of substances affecting reticuloendothelial system on level of, mechanism of action of Riley virus, 983
Virus, Riley, mechanism of action of substances affecting reticuloendothelial system on level of viremia, 983
—, Riley, mechanism of action of substances affecting reticuloendothelial system on plasma enzyme levels, 967
—, Riley, mechanism of action, replication of Riley's plasma enzyme-elevating virus *in vitro*, 993
Viruses, adenovirus SV40 hybrid, assay system and further evidence for hybridization, 943
—, adenovirus SV40 hybrid, defectiveness of hybrid particles, 955

WALL, cell, mucopeptide, enhanced resistance to streptococcal infection induced in mice by, 877